





تبادل اسید-باز (قسمت سوم)

یونش خود به خودی آب و مقیاس pH:



$$K_c = \frac{[\text{H}_3\text{O}^+][\text{OH}^-]}{[\text{H}_2\text{O}]^2}$$

The Ion-Product Constant for Water

$$K_c[\text{H}_2\text{O}]^2 = K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1.0 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

A change in $[\text{H}_3\text{O}^+]$ causes an inverse change in $[\text{OH}^-]$.

In an acidic solution, $[\text{H}_3\text{O}^+] > [\text{OH}^-]$

In a basic solution, $[\text{H}_3\text{O}^+] < [\text{OH}^-]$

In a neutral solution, $[\text{H}_3\text{O}^+] = [\text{OH}^-]$

$$[\text{H}_3\text{O}^+] > [\text{OH}^-]$$



**ACIDIC
SOLUTION**

$$[\text{H}_3\text{O}^+] = [\text{OH}^-]$$



**NEUTRAL
SOLUTION**

$$[\text{H}_3\text{O}^+] < [\text{OH}^-]$$



**BASIC
SOLUTION**

Sample Problem 18.2

Calculating $[H_3O^+]$ and $[OH^-]$ in an Aqueous Solution

PROBLEM: A research chemist adds a measured amount of HCl gas to pure water at 25°C and obtains a solution with $[H_3O^+] = 3.0 \times 10^{-4} M$. Calculate $[OH^-]$. Is the solution neutral, acidic, or basic?

PLAN: Use the K_w at 25°C and the $[H_3O^+]$ to find the corresponding $[OH^-]$.

SOLUTION: $K_w = 1.0 \times 10^{-14} = [H_3O^+] [OH^-]$ so


$$[OH^-] = K_w / [H_3O^+] = 1.0 \times 10^{-14} / 3.0 \times 10^{-4} = 3.3 \times 10^{-11} M$$

$[H_3O^+]$ is $>$ $[OH^-]$ and the solution is acidic.

تعریف pH و مقیاس آن

Figure 18.5
The pH values of some familiar aqueous solutions.

$$pH = -\log [H_3O^+]$$



محاسبه غلظت یونهای هیدرونیوم و هیدروکسید و تابع p آنها در محلول اسید قوی

به عنوان مثال برای محلولهای اسیدی که برای کار روی صفحات مسی در آثار هنری

به کار می رود:

Sample Problem 18.3 Calculating $[H_3O^+]$, pH, $[OH^-]$, and pOH


PROBLEM: In an art restoration project, a conservator prepares copper-plate etching solutions by diluting concentrated HNO_3 to 2.0 M, 0.30 M, and 0.0063 M HNO_3 . Calculate $[H_3O^+]$, pH, $[OH^-]$, and pOH of the three solutions at 25°C.

PLAN: HNO_3 is a strong acid so $[H_3O^+] = [HNO_3]$. Use K_w to find the $[OH^-]$ and then convert to pH and pOH.

SOLUTION: For 2.0 M HNO_3 , $[H_3O^+] = 2.0 M$ and $-\log [H_3O^+] = -0.30 = \text{pH}$
 $[OH^-] = K_w / [H_3O^+] = 1.0 \times 10^{-14} / 2.0 = 5.0 \times 10^{-15} M$; pOH = 14.30

For 0.30 M HNO_3 , $[H_3O^+] = 0.30 M$ and $-\log [H_3O^+] = 0.52 = \text{pH}$
 $[OH^-] = K_w / [H_3O^+] = 1.0 \times 10^{-14} / 0.30 = 3.3 \times 10^{-14} M$; pOH = 13.48


For 0.0063 M HNO_3 , $[H_3O^+] = 0.0063 M$ and $-\log [H_3O^+] = 2.20 = \text{pH}$
 $[OH^-] = K_w / [H_3O^+] = 1.0 \times 10^{-14} / 6.3 \times 10^{-3} = 1.6 \times 10^{-12} M$; pOH = 11.80



محاسبه pH محلول باز قوی


تمرین:

محاسبه pH محلولی از سود با غلظت 0/02 مولار



اسیدهای ضعیف

- اسید ضعیف مانند استیک اسید
- اسید مزدوج بازهای ضعیف مانند یون آمونیوم



$$\text{Percent HA dissociation} = \frac{[\text{HA}]_{\text{dissociated}}}{[\text{HA}]_{\text{initial}}} \times 100$$

$$\% \text{ dissociated} = 4/(5 + 4) \times 100 = 44\%$$

دو رویکرد در مسائل مربوط به تعادل اسید ضعیف:

- یافتن K_a از pH محلول

نمونه مسئله 7-18: یافتن K_a محلول 0/12 مولار فنیل استیک اسید با pH = 2.62.

Sample Problem 18.7

Finding the K_a of a Weak Acid from the Solution pH

PROBLEM: Phenylacetic acid ($\text{C}_6\text{H}_5\text{CH}_2\text{COOH}$, simplified here as HPAC) builds up in the blood of persons with phenylketonuria, an inherited disorder that, if untreated, causes mental retardation and death. A study of the acid shows that the pH of 0.12 M HPAC is 2.62. What is the K_a of phenylacetic acid?

Concentration (M)	HPAC(aq) +	$\text{H}_2\text{O}(l)$	\rightleftharpoons	$\text{H}_3\text{O}^+(aq)$ +	PAC ⁻ (aq)
Initial	0.12	-		0	0
Change	-x	-		+x	+x
Equilibrium	0.12 - x	-		x	x

$$[\text{H}_3\text{O}^+] = 10^{-\text{pH}} = 2.4 \times 10^{-3} \text{ M which is } \gg 10^{-7} \text{ (the } [\text{H}_3\text{O}^+] \text{ from water)}$$

$$x \approx 2.4 \times 10^{-3} \text{ M} \approx [\text{H}_3\text{O}^+] \approx [\text{PAC}^-] \quad [\text{HPAC}]_{\text{equilibrium}} = 0.12 - x \approx 0.12 \text{ M}$$

$$\text{So } K_a = \frac{(2.4 \times 10^{-3})(2.4 \times 10^{-3})}{0.12} = 4.8 \times 10^{-5}$$

- یافتن غلظت‌های تعادلی با داشتن K_a و غلظت اولیه اسید
 نمونه مسئله 8-18 پروپانویک اسید:

Sample Problem 18.8

Determining Concentrations from K_a and Initial [HA]

PROBLEM: Propanoic acid ($\text{CH}_3\text{CH}_2\text{COOH}$, which we simplify as HPr) is a organic acid whose salts are used to retard mold growth in foods. What is the $[\text{H}_3\text{O}^+]$ of 0.10 M HPr ($K_a = 1.3 \times 10^{-5}$)?

PLAN: Write out the dissociation equation and expression; make whatever assumptions about concentration which are necessary; substitute.

Assumptions: For $\text{HPr}(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{Pr}^-(aq)$

$$x = [\text{HPr}]_{\text{diss}} = [\text{H}_3\text{O}^+]_{\text{from HPr}} = [\text{Pr}^-] \quad K_a = \frac{[\text{H}_3\text{O}^+][\text{Pr}^-]}{[\text{HPr}]}$$

SOLUTION:

Concentration (M)	HPr(aq)	+ H ₂ O(l)	\rightleftharpoons	H ₃ O ⁺ (aq)	+ Pr ⁻ (aq)
Initial	0.10	-		0	0
Change	-x	-		+x	+x
Equilibrium	0.10 - x	-		x	x

Since K_a is small, we will assume that $x \ll 0.10$

$$1.3 \times 10^{-5} = \frac{[\text{H}_3\text{O}^+][\text{Pr}^-]}{[\text{HPr}]} = \frac{(x)(x)}{0.10} \quad \boxed{\text{تقریب}}$$

$$x = \sqrt{(0.10)(1.3 \times 10^{-5})} = 1.1 \times 10^{-3} \text{ M} = [\text{H}_3\text{O}^+]$$




$$\text{Check: } [\text{HPr}]_{\text{diss}} = 1.1 \times 10^{-3} \text{ M} / 0.10 \text{ M} \times 100 = 1.1\%$$

اسیدهای چند پروتونی:

در این اسیدها هر مرحله یونش در pH موثر است ولی اگر ثابت‌های تعادل دوم و سوم خیلی کوچکتر از ثابت تفکیک اول باشد از مراحل بعدی صرفنظر می‌شود.

Polyprotic acids

acids with more than one ionizable proton



$$K_{a1} > K_{a2} > K_{a3}$$

نمونه مسئله 18-10: تعیین غلظتهای تعادلی در محلول آسکوربیک اسید 0/050 مولار (اسید دو ظرفیتی)

Sample Problem 18.10

Calculating Equilibrium Concentrations for a Polyprotic Acid

PROBLEM: Ascorbic acid ($\text{H}_2\text{C}_6\text{H}_6\text{O}_6$; H_2Asc for this problem), known as vitamin C, is a diprotic acid ($K_{a1} = 1.0 \times 10^{-5}$ and $K_{a2} = 5 \times 10^{-12}$) found in citrus fruit. Calculate $[\text{H}_2\text{Asc}]$, $[\text{HAsc}^-]$, $[\text{Asc}^{2-}]$, and the pH of 0.050 M H_2Asc .



PLAN: Write out expressions for both dissociations and make assumptions.

$K_{a1} \gg K_{a2}$ so the first dissociation produces virtually all of the H_3O^+ .

K_{a1} is small so $[\text{H}_2\text{Asc}]_{\text{initial}} \approx [\text{H}_2\text{Asc}]$


After finding the concentrations of various species for the first dissociation, we can use them as initial concentrations for the second dissociation.

SOLUTION:



Sample Problem 18.10

Calculating Equilibrium Concentrations for a Polyprotic Acid




Initial	0.050	-	0	0
Change	-x	-	+x	+x
Equilibrium	0.050 - x	-	x	x

$$K_{a1} = \frac{[\text{HAsc}^-][\text{H}_3\text{O}^+]}{[\text{H}_2\text{Asc}]} = 1.0 \times 10^{-5} = \frac{(x)(x)}{0.050} \text{ M}$$

تقریب

$$x = \sqrt{(0.050)(1.0 \times 10^{-5})} \quad x = 7.1 \times 10^{-4} \text{ M}$$

$$\text{pH} = -\log(7.1 \times 10^{-4}) = 3.15$$



Initial	$7.1 \times 10^{-4} \text{ M}$	-	0	$7.1 \times 10^{-4} \text{ M}$
Change	-x	-	+x	+x
Equilibrium	$7.1 \times 10^{-4} - x$	-	x	$7.1 \times 10^{-4} + x$

$$K_{a2} = \frac{[\text{Asc}^{2-}][\text{H}_3\text{O}^+]}{[\text{HAsc}^-]} = 5 \times 10^{-12} = \frac{(x)(7.1 \times 10^{-4} + x)}{(7.1 \times 10^{-4} - x)} \text{ M} = x$$

18-35

تقریب بزنیید

اصلاح کنید